

**Table 4. Estimated distribution of reported results of the first CD4 test performed within 12 months after HIV diagnosis among adults and adolescents, by disease category and sex, 2001–2003—33 states with confidential name-based HIV infection reporting**

	CD4 count, cells/μL <sup>a</sup>				No CD4 count	Total
	<200 No. (%)	200–350 No. (%)	351–500 No. (%)	>500 No. (%)		
HIV without AIDS <sup>c</sup>						
Males	— —	4,348 (8.7)	4,998 (10.0)	6,724 (13.4)	34,023 (67.9)	50,094
Females	— —	1,855 (8.1)	2,015 (8.8)	2,996 (13.1)	15,975 (69.9)	22,840
Subtotal	— —	6,203 (8.5)	7,013 (9.6)	9,720 (13.3)	49,997 (68.6)	72,933
HIV to AIDS <sup>d</sup>						
Males	11,129 (80.1)	1,572 (11.3)	514 (3.7)	271 (2.0)	405 (2.9)	13,890
Females	4,210 (81.6)	546 (10.6)	181 (3.5)	92 (1.8)	129 (2.5)	5,157
Subtotal	15,338 (80.5)	2,118 (11.1)	694 (3.6)	363 (1.9)	533 (2.8)	19,047
HIV with AIDS <sup>e</sup>						
Males	16,618 (85.7)	1,016 (5.2)	326 (1.7)	196 (1.0)	1,233 (6.4)	19,388
Females	5,332 (86.1)	296 (4.8)	89 (1.4)	71 (1.1)	406 (6.6)	6,195
Subtotal	21,951 (85.8)	1,312 (5.1)	415 (1.6)	267 (1.0)	1,639 (6.4)	25,583

*Note.* These numbers do not represent actual cases in persons with a diagnosis of HIV infection or AIDS. Rather, these numbers are point estimates of case counts that have been adjusted for reporting delays. The estimates have not been adjusted for incomplete reporting.

Since 2000, the following states have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Alaska, Arizona, Arkansas, Colorado, Florida, Idaho, Indiana, Iowa, Kansas, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, West Virginia, Wisconsin, and Wyoming. Since July 1997, Florida has had confidential name-based HIV infection reporting for new diagnoses only.

Dash indicates not applicable.

<sup>a</sup> When only CD4 percentage was available, a CD4 count was interpreted.

<sup>b</sup> Because row totals were calculated independently of the row values, the values in each row may not sum to the row total.

<sup>c</sup> For at least 12 months after HIV diagnosis.

<sup>d</sup> AIDS diagnosed 1–12 months after HIV diagnosis.

<sup>e</sup> HIV and AIDS diagnosed during the same calendar month.